



Ferinject Infusion Patient Consent Form

Informed Consent to Receive Intravenous Iron Replacement Therapy

The patient understands that the administration of FERINJECT comes with the following risks, included but not limited to:

- Anaphylactic reactions, which in rare cases may be potentially fatal
 - Paravenous leakage- leakage in FERINJECT at injection site, potentially leading to long lasting skin discolouration
 - Skin Irritations
 - Headaches, light headiness
 - Tachycardia, Hyper/Hypotension
 - Nausea, Stomach pain, Constipation, Diarrhoea and Vomiting
- Minor reactions to FERINJECT may last up to 48 hours post injection. For full list of reactions, please see page 2.

Understanding these risks, patient gives authority for staff of Mooroolbark Medical Centre to administer all necessary first aid and/or resuscitation measures, including alerting an ambulance and my Emergency Contact, in the unlikely event that an adverse or anaphylactic reaction occurs.

As FERINJECT is **not suitable** for patients in some conditions, patient declares that none of the below listed is applicable:

- Pregnancy in the first trimester
- Dialysis
- Allergy to Ferric Carboxymaltose
- Iron overload/Haemochromotosis
- Under the age of 14 years
- Non-iron deficiency related anaemia
- Suffering from fever/sepsis

The patient, as stated below, has read information provided on this document as well as Mooroolbark Medical Centre's Policy and Procedures relating to the administration of FERINJECT.

The patient understands that procedure is undertaken entirely at own risk and cost and is requesting medical intervention in the form of Intravenous Iron Infusion.

Patient Full Name: DOB:

Patient Full Address:

Patient Signature: Date:

Doctor Signature: Date:

Copy of Blood Test obtained and scanned to patient file

Ferinject Infusion Patient Consent Form



Common Side Effects (Occurs in 1-10% of cases):

- Headache
- Flushing
- Nausea
- Low blood phosphate levels
- Injection/Infusion site reactions

Uncommon and Rare Side Effects (Occurs in <1% of cases):

- Hypersensitivity
- Numbness
- Increased heart rate
- Hypotension
- Difficulty breathing
- Taste disturbance
- Vomiting
- Dyspepsia
- Flatulence
- Abdominal pain
- Constipation
- Diarrhoea
- Itchiness
- Hives (urticaria)
- Redness (erythema)
- Rash
- Muscle pain or spasm
- Back, joint or chest pain
- Fever
- Fatigue
- Accumulation of fluid in periphery
- Pain and chills
- Anaphylaxis
- Rigors
- Malaise